

Psychosis and schizophrenia in children and young people

Recognition and management

Issued: January 2013

NICE clinical guideline 155

guidance.nice.org.uk/cg155

Contents

Introduction	3
Person-centred care	6
Key priorities for implementation.....	7
1 Recommendations	11
1.1 General principles of care	11
1.2 Possible psychosis.....	17
1.3 First episode psychosis.....	18
1.4 Subsequent acute episodes of psychosis or schizophrenia	27
1.5 Referral in crisis and challenging behaviour.....	29
1.6 Early post-acute period	32
1.7 Promoting recovery and providing possible future care in primary care.....	32
1.8 Promoting recovery and providing possible future care in secondary care	34
2 Research recommendations.....	37
3 Other information.....	41
3.1 Scope and how this guideline was developed.....	41
3.2 Related NICE guidance.....	42
4 The Guideline Development Group, National Collaborating Centre and NICE project team	43
Supplementary information on baseline investigations and monitoring.....	47
Table 1 Baseline investigations and monitoring for children and young people who are prescribed antipsychotic medication (read in conjunction with the BNF, BNFC and SPC).	47
About this guideline.....	48
Strength of recommendations	49
Other versions of this guideline	50
Implementation.....	50
Your responsibility.....	50
Copyright.....	51
Contact NICE	51

Introduction

This guideline is concerned with the recognition and management of psychosis and schizophrenia in children and young people up to the age of 18. The term 'psychosis' is used in this guideline to refer to the group of psychotic disorders that includes schizophrenia, schizoaffective disorder, schizophreniform disorder and delusional disorder. This guideline also addresses those children and young people considered clinically to be at high risk or prodromal for psychosis and schizophrenia. The recognition, treatment and management of affective psychoses (such as bipolar disorder or unipolar psychotic depression) are covered by other NICE guidelines.

Psychosis and the specific diagnosis of schizophrenia in children and young people represent a major psychiatric disorder, or cluster of disorders that alters a person's perception, thoughts, mood and behaviour. The symptoms of psychosis are usually divided into 'positive symptoms', including hallucinations (perception in the absence of any stimulus) and delusions (fixed or falsely held beliefs), and 'negative symptoms' (such as emotional apathy, lack of drive, poverty of speech, social withdrawal and self-neglect). Children and young people who develop psychosis will have their own unique combination of symptoms and experiences, the precise pattern of which will be influenced by their circumstances and stage of development.

Psychosis and schizophrenia are commonly preceded by a so-called prodromal period, lasting up to 12 months, in which the child or young person's behaviour and experience are altered. Relatives may become aware of these changes first. Changes include the emergence of transient and/or attenuated psychotic symptoms, such as hallucinations and/or delusions with associated impaired functioning. More subtly, the child or young person may become socially withdrawn or suspicious, with alterations in expressed feeling. It is important to note that most children and young people with transient or attenuated psychotic symptoms do not go on to develop psychosis or schizophrenia, although those with such symptoms do appear to be at higher risk than other children and young people of developing psychosis and schizophrenia up to 10 years after onset of symptoms.

The prevalence of psychotic disorders in children aged between 5 and 18 years has been estimated to be 0.4% (the figure across all ages and populations in the UK is 0.7%). Schizophrenia accounts for 24.5% of all psychiatric admissions in young people aged 10–

18 years (the overall admission rate is 0.46 per 1000 for this age range), with an exponential rise across the adolescent years. The rise in incidence increases most from age 15 onwards.

There is a worse prognosis for psychosis and schizophrenia when onset is in childhood or adolescence. The symptoms and experience of psychosis and schizophrenia are often distressing and the effects of the illness are pervasive. Although about one-fifth of children and young people with schizophrenia have a good outcome with only mild impairment, one-third have severe impairment that needs intensive social and psychiatric support. Psychosis and schizophrenia can have a major detrimental effect on children and young people's personal, social, educational and occupational functioning, placing a heavy burden on them and their parents and carers.

Although the mainstay of treatment for psychosis and schizophrenia has been antipsychotic medication, there is limited evidence of its efficacy in children and young people. There are also concerns that children and young people are more sensitive than adults to the potential adverse effects of antipsychotics, including weight gain, metabolic effects and movement disorders. A number of psychological interventions, including family intervention, cognitive behavioural therapy (CBT) and arts therapies, have been used but evidence of efficacy is currently unavailable in children and young people and provision of these therapies for children and young people and for adults is variable.

This guideline covers the care provided by primary, community, secondary, tertiary and other health and social care professionals who have direct contact with, and make decisions concerning, the care of children and young people with psychosis or schizophrenia, including child and adolescent mental health services (CAMHS) and early intervention in psychosis services.

Early intervention in psychosis services provide people aged 14–35 years with a more intensive therapeutic service than traditional community services. They are designed to intervene early, and deliver support and evidence-based interventions in a 'normalising' environment for the first 3 years after onset of psychosis.

There is geographical variation in the configuration and integration of CAMHS and early intervention in psychosis services, and in the provision and integration of other services for children and young people with psychosis and schizophrenia, including education, employment

and rehabilitation, and social services. In particular, provision for the needs of 16- and 17-year-olds with psychosis and schizophrenia can be fragmented and inadequate and they can experience difficulties in gaining access to appropriate accommodation and vocational or occupational support and rehabilitation.

A number of recommendations in this guideline have been adapted from recommendations in other NICE clinical guidelines. Where this occurred, the Guideline Development Group was careful to preserve the meaning and intent of the original recommendation. Changes to wording or structure were made in order to fit the recommendations into this guideline. In all cases, the original source of any adapted recommendation is indicated in a footnote.

The guideline incorporates [Aripiprazole for the treatment of schizophrenia in people aged 15 to 17 years](#) (NICE technology appraisal guidance 213).

The guideline will assume that prescribers will use a drug's summary of product characteristics (SPC) to inform decisions made with individual patients.

This guideline recommends some drugs for indications for which they do not have a UK marketing authorisation at the date of publication, if there is good evidence to support that use. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. The patient (or those with authority to give consent on their behalf) should provide informed consent, which should be documented. See the General Medical Council's [Good practice in prescribing medicines – guidance for doctors](#) for further information. Where recommendations have been made for the use of drugs outside their licensed indications ('off-label use'), these drugs are marked with a footnote in the recommendations.

Person-centred care

This guideline offers best practice advice on the care of children and young people with psychosis or schizophrenia.

Service users and healthcare professionals have rights and responsibilities as set out in the [NHS Constitution for England](#) – all NICE guidance is written to reflect these. Treatment and care should take into account individual needs and preferences. Service users should have the opportunity to make informed decisions about their care and treatment, in partnership with their healthcare professionals. If someone does not have the capacity to make decisions, healthcare professionals should follow the [Department of Health's advice on consent](#) and the [code of practice that accompanies the Mental Capacity Act](#) and the supplementary [code of practice on deprivation of liberty safeguards](#). In Wales, healthcare professionals should follow [advice on consent from the Welsh Government](#).

If the service user is under 16, healthcare professionals should follow the guidelines in the Department of Health's [Seeking consent: working with children](#). Parents and carers should also be given the information and support they need to help the child or young person in making decisions about their treatment.

If a young person is moving between CAMHS and adult mental health services, care should be planned and managed according to the best practice guidance described in the Department of Health's [Transition: getting it right for young people](#).

CAMHS and adult mental health services should work jointly to provide assessment and services to children and young people with psychosis or schizophrenia. Diagnosis and management should be reviewed throughout the transition process, and there should be clarity about who is the lead clinician to ensure continuity of care.

Key priorities for implementation

The following recommendations have been identified as priorities for implementation.

Referral from primary care for possible psychosis

- When a child or young person experiences transient or attenuated psychotic symptoms or other experiences suggestive of possible psychosis, refer for assessment without delay to a specialist mental health service such as CAMHS or an early intervention in psychosis service (14 years or over).

Treatment options for symptoms not sufficient for a diagnosis of psychosis or schizophrenia

- When transient or attenuated psychotic symptoms or other mental state changes associated with distress, impairment or help-seeking behaviour are not sufficient for a diagnosis of psychosis or schizophrenia:
 - consider individual cognitive behavioural therapy (CBT) (delivered as set out in recommendation 1.3.28) with or without family intervention (delivered as set out in recommendation 1.3.27), and
 - offer treatments recommended in NICE guidance for children and young people with any of the anxiety disorders, depression, emerging personality disorder or substance misuse.
- Do not offer antipsychotic medication:
 - for psychotic symptoms or mental state changes that are not sufficient for a diagnosis of psychosis or schizophrenia, or
 - with the aim of decreasing the risk of psychosis.

Referral from primary care for first episode psychosis

- Urgently refer all children and young people with a first presentation of sustained psychotic symptoms (lasting 4 weeks or more) to a specialist mental health service, either CAMHS (up to 17 years) or an early intervention in psychosis service (14 years or over), which includes a consultant psychiatrist with training in child and adolescent mental health.

Treatment options for first episode psychosis

- If the child or young person and their parents or carers wish to try psychological interventions (family intervention with individual CBT) alone without antipsychotic medication, advise that psychological interventions are more effective when delivered in conjunction with antipsychotic medication. If the child or young person and their parents or carers still wish to try psychological interventions alone, then offer family intervention with individual CBT. Agree a time limit (1 month or less) for reviewing treatment options, including introducing antipsychotic medication. Continue to monitor symptoms, level of distress, impairment and level of functioning, including educational engagement and achievement, regularly.

How to use oral antipsychotic medication

- Before starting antipsychotic medication¹, undertake and record the following baseline investigations²:
 - weight and height (both plotted on a growth chart)
 - waist and hip circumference
 - pulse and blood pressure
 - fasting blood glucose, glycosylated haemoglobin (HbA_{1c}), blood lipid profile and prolactin levels
 - assessment of any movement disorders
 - assessment of nutritional status, diet and level of physical activity.
- Monitor and record the following regularly and systematically throughout treatment, but especially during titration²:
 - efficacy, including changes in symptoms and behaviour
 - side effects of treatment, taking into account overlap between certain side effects and clinical features of schizophrenia (for example, the overlap between akathisia and agitation or anxiety)
 - the emergence of movement disorders

¹ At the time of publication (January 2013), most antipsychotic medication did not have a UK marketing authorisation specifically for children and young people. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. Informed consent should be obtained and documented. See the General Medical Council's [Good practice in prescribing medicines – guidance for doctors](#) for further information.

² See [Supplementary information](#) for a table of baseline investigations and monitoring for children and young people who are prescribed antipsychotic medication (read in conjunction with the BNF, BNFC and SPC).

-
- weight, weekly for the first 6 weeks, then at 12 weeks and then every 6 months (plotted on a growth chart)
 - height every 6 months (plotted on a growth chart)
 - waist and hip circumference every 6 months (plotted on a percentile chart)
 - pulse and blood pressure (plotted on a percentile chart) at 12 weeks and then every 6 months
 - fasting blood glucose, HbA_{1c}, blood lipid and prolactin levels at 12 weeks and then every 6 months
 - adherence
 - physical health.

The secondary care team should maintain responsibility for monitoring physical health and the effects of antipsychotic medication in children and young people for at least the first 12 months or until their condition has stabilised. Thereafter, the responsibility for this monitoring may be transferred to primary care under shared care arrangements.

Treatment of subsequent acute episodes of psychosis or schizophrenia

- Offer family intervention (delivered as set out in recommendation 1.3.27) to all families of children and young people with psychosis or schizophrenia, particularly for preventing and reducing relapse. This can be started either during the acute phase or later, including in inpatient settings³.
- Before referral for hospital care, think about the impact on the child or young person and their parents, carers and other family members, especially when the inpatient unit is a long way from where they live. Consider alternative care within the community wherever possible. If hospital admission is unavoidable, provide support for parents or carers when the child or young person is admitted.

³ Adapted from [Schizophrenia](#) (NICE clinical guideline 82).

Education, employment and occupational activities for children and young people with psychosis and schizophrenia

- For children and young people of compulsory school age, liaise with the child or young person's school and educational authority, subject to consent, to ensure that ongoing education is provided.

1 Recommendations

The following guidance is based on the best available evidence. The [full guideline](#) gives details of the methods and the evidence used to develop the guidance.

All recommendations relate to children and young people (younger than 18 years) unless otherwise specified.

1.1 *General principles of care*

Working safely and effectively with children and young people

- 1.1.1 Health and social care professionals working with children and young people with psychosis or schizophrenia should be trained and competent to work with children and young people with mental health problems of all levels of learning ability, cognitive capacity, emotional maturity and development.
- 1.1.2 Health and social care professionals should ensure that they:
- can assess capacity and competence, including 'Gillick competence', in children and young people of all ages, and
 - understand how to apply legislation, including the Children Act (1989; amended 2004), the Mental Health Act (1983; amended 1995 and 2007⁴) and the Mental Capacity Act (2005), in the care and treatment of children and young people.
- 1.1.3 Consider children and young people with psychosis or schizophrenia for assessment according to local safeguarding procedures if there are concerns regarding exploitation or self-care, or if they have been in contact with the criminal justice system⁵.
- 1.1.4 Health and social care providers should ensure that children and young people with psychosis or schizophrenia:
- can routinely receive care and treatment from a single multidisciplinary community team

⁴ Including the [Code of Practice: Mental Health Act 1983](#).

⁵ Adapted from [Service user experience in adult mental health](#) (NICE clinical guidance 136).

- are not passed from one team to another unnecessarily
- do not undergo multiple assessments unnecessarily⁵.

1.1.5 Help the child or young person to continue their education. Contact the school or college, subject to consent, to ask for additional educational support if their performance has been affected by their condition.

Establishing relationships with children and young people and their parents or carers

1.1.6 Work in partnership with children and young people with psychosis or schizophrenia of an appropriate developmental level, emotional maturity and cognitive capacity and parents or carers. Offer help, treatment and care in an atmosphere of hope and optimism. Take time to build trusting, supportive, empathic and non-judgemental relationships as an essential part of care⁵.

1.1.7 When working with children and young people with psychosis or schizophrenia:

- aim to foster autonomy, promote active participation in treatment decisions, and support self-management and access to peer support in children and young people of an appropriate developmental level, emotional maturity and cognitive capacity
- maintain continuity of individual therapeutic relationships wherever possible
- offer access to a trained advocate⁵.

1.1.8 When working with children and young people with psychosis or schizophrenia and their parents or carers:

- make sure that discussions take place in settings in which confidentiality, privacy and dignity are respected
- be clear with the child or young person and their parents or carers about limits of confidentiality (that is, which health and social care professionals have access to information about their diagnosis and its treatment and in what circumstances this may be shared with others)⁵.

- 1.1.9 Discuss with young people with psychosis or schizophrenia of an appropriate developmental level, emotional maturity and cognitive capacity how they want their parents or carers to be involved in their care. Such discussions should take place at intervals to take account of any changes in circumstances, including developmental level, and should not happen only once⁵.
- 1.1.10 Advise parents and carers about their right to a formal carer's assessment of their own physical and mental health needs, and explain how to access this⁵.

Communication and information

- 1.1.11 Health and social care professionals working with children and young people with psychosis or schizophrenia should be trained and skilled in:
- negotiating and working with parents and carers, and
 - managing issues relating to information sharing and confidentiality as these apply to children and young people.
- 1.1.12 If a young person is 'Gillick competent' ask them what information can be shared before discussing their condition and treatment with their parents or carers.
- 1.1.13 When communicating with children and young people with psychosis or schizophrenia and their parents or carers:
- take into account the child or young person's developmental level, emotional maturity and cognitive capacity including any learning disabilities, sight or hearing problems or delays in language development
 - use plain language where possible and clearly explain any clinical language
 - check that the child or young person and their parents or carers understand what is being said
 - use communication aids (such as pictures, symbols, large print, braille, different languages or sign language) if needed.

- 1.1.14 Provide children and young people with psychosis or schizophrenia and their parents or carers, comprehensive written information about:
- the nature of, and interventions for, psychosis and schizophrenia (including biomedical and psychosocial perspectives on causes and treatment) in an appropriate language or format, including any relevant 'Information for the public' booklets
 - support groups, such as third sector, including voluntary, organisations⁵.
- 1.1.15 Ensure that you are:
- familiar with local and national sources (organisations and websites) of information and/or support for children and young people with psychosis or schizophrenia and their parents or carers
 - able to discuss and advise how to access these resources
 - able to discuss and actively support children and young people and their parents or carers to engage with these resources⁵.
- 1.1.16 When communicating with a child or young person with psychosis or schizophrenia, use diverse media, including letters, phone calls, emails or text messages, according to their preference⁵.
- 1.1.17 Copy all written communications with other health or social care professionals to the child or young person and/or their parents or carers at the address of their choice, unless this is declined⁵.

Culture, ethnicity and social inclusion

- 1.1.18 When working with children and young people with psychosis or schizophrenia and their parents or carers:
- take into account that stigma and discrimination are often associated with using mental health services
 - be respectful of and sensitive to children and young people's gender, sexual orientation, socioeconomic status, age, background (including cultural, ethnic and religious background) and any disability

- be aware of possible variations in the presentation of mental health problems in children and young people of different genders, ages, cultural, ethnic, religious or other diverse backgrounds⁵.

1.1.19 When working with children and young people and their parents or carers who have difficulties speaking or reading English:

- provide and work proficiently with interpreters if needed
- offer a list of local education providers who can provide English language teaching.

1.1.20 Health and social care professionals working with children and young people with psychosis or schizophrenia and their parents or carers should have competence in:

- assessment skills for people from diverse ethnic and cultural backgrounds
- using explanatory models of illness for people from diverse ethnic and cultural backgrounds
- explaining the possible causes of psychosis and schizophrenia and treatment options
- addressing cultural and ethnic differences in treatment expectations and adherence
- addressing cultural and ethnic differences in beliefs regarding biological, social and family influences on the possible causes of mental health problems
- conflict management and conflict resolution⁶.

1.1.21 Health and social care professionals inexperienced in working with children and young people with psychosis or schizophrenia from diverse ethnic and cultural backgrounds, and their parents or carers, should seek advice and supervision from healthcare professionals who are experienced in working transculturally⁶.

⁶ Adapted from [Schizophrenia](#) (NICE clinical guideline 82).

- 1.1.22 Local mental health services should work with primary care, other secondary care and local third sector, including voluntary, organisations to ensure that:
- all children and young people with psychosis or schizophrenia have equal access to services based on clinical need and irrespective of gender, sexual orientation, socioeconomic status, age, background (including cultural, ethnic and religious background) and any disability
 - services are culturally appropriate⁵.
- 1.1.23 Mental health services should work with local voluntary black and minority ethnic groups to jointly ensure that culturally appropriate psychological and psychosocial treatment, consistent with this guideline and delivered by competent practitioners, is provided to children and young people from diverse ethnic and cultural backgrounds⁶.

Transfer and discharge

- 1.1.24 Anticipate that withdrawal and ending of treatments or services, and transition from one service to another, may evoke strong emotions and reactions in children and young people with psychosis or schizophrenia and their parents or carers. Ensure that:
- such changes, especially discharge and transfer from CAMHS to adult services, or to primary care, are discussed and planned carefully beforehand with the child or young person and their parents or carers, and are structured and phased
 - the care plan supports effective collaboration with social care and other care providers during endings and transitions, and includes details of how to access services in times of crisis
 - when referring a child or young person for an assessment in other services (including for psychological interventions), they are supported during the referral period and arrangements for support are agreed beforehand with them⁵.

1.2 *Possible psychosis*

Referral from primary care

- 1.2.1 When a child or young person experiences transient or attenuated psychotic symptoms or other experiences suggestive of possible psychosis, refer for assessment without delay to a specialist mental health service such as CAMHS or an early intervention in psychosis service (14 years or over).

Assessment in specialist mental health services

- 1.2.2 Carry out an assessment of the child or young person with possible psychosis, ensuring that:
- assessments in CAMHS include a consultant psychiatrist
 - assessments in early intervention in psychosis services are multidisciplinary
 - where there is considerable uncertainty about the diagnosis, or concern about underlying neurological illness, there is an assessment by a consultant psychiatrist with training in child and adolescent mental health.
- 1.2.3 If a clear diagnosis of psychosis cannot be made, monitor regularly for further changes in symptoms and functioning for up to 3 years. Determine the frequency and duration of monitoring by:
- the severity and frequency of symptoms
 - the level of impairment and/or distress in the child or young person, and
 - the degree of family disruption or concern.
- 1.2.4 If discharge from the service is requested, offer follow-up appointments and the option to self-refer at a later date. Ask the GP to continue monitoring changes in mental state.

Treatment options for symptoms not sufficient for a diagnosis of psychosis or schizophrenia

1.2.5 When transient or attenuated psychotic symptoms or other mental state changes associated with distress, impairment or help-seeking behaviour are not sufficient for a diagnosis of psychosis or schizophrenia:

- consider individual cognitive behavioural therapy (CBT) (delivered as set out in recommendation 1.3.28) with or without family intervention (delivered as set out in recommendation 1.3.27), and
- offer treatments recommended in NICE guidance for children and young people with any of the anxiety disorders, depression, emerging personality disorder or substance misuse.

1.2.6 Do not offer antipsychotic medication:

- for psychotic symptoms or mental state changes that are not sufficient for a diagnosis of psychosis or schizophrenia, or
- with the aim of decreasing the risk of psychosis.

1.3 *First episode psychosis*

Referral from primary care

1.3.1 Urgently refer all children and young people with a first presentation of sustained psychotic symptoms (lasting 4 weeks or more) to a specialist mental health service, either CAMHS (up to 17 years) or an early intervention in psychosis service (14 years or over), which includes a consultant psychiatrist with training in child and adolescent mental health.

1.3.2 Antipsychotic medication in children and young people with a first presentation of sustained psychotic symptoms should not be started in primary care unless it is done in consultation with a consultant psychiatrist with training in child and adolescent mental health.

Assessment and care planning in secondary care

1.3.3 When carrying out an assessment:

- ensure there is enough time for:
 - the child or young person and their parents or carers to describe and discuss their problems
 - summarising the conclusions of the assessment and for discussion, with questions and answers
- explain and give written material in an accessible format about any diagnosis given
- give information about different treatment options, including pharmacological and psychological interventions, and their benefits and side effects, to promote discussion and shared understanding
- offer support after the assessment, particularly if sensitive issues, such as childhood trauma, have been discussed⁵.

1.3.4 Ensure that children and young people with first episode psychosis receive a comprehensive multidisciplinary assessment. The assessment should address the following domains:

- psychiatric (mental health problems, risk of harm to self or others, alcohol consumption and prescribed and non-prescribed drug history)
- medical, including medical history and full physical examination to identify physical illness (including organic brain disorders) and prescribed drug treatments that may result in psychosis
- psychological and psychosocial, including social networks, relationships and history of trauma
- developmental (social, cognitive and motor development and skills, including coexisting neurodevelopmental conditions)
- physical health and wellbeing (including weight and height, and information about smoking, diet and exercise, and sexual health)
- social (accommodation, culture and ethnicity, leisure activities and recreation, carer responsibilities [for example, of parents or siblings])

- educational and occupational (attendance at school or college, educational attainment, employment and functional activity)
- economic (family's economic status).

1.3.5 Routinely monitor for other coexisting mental health problems, including depression and anxiety, and substance misuse, particularly in the early phases of treatment⁶.

1.3.6 Develop a care plan with the parents or carers of younger children, or jointly with the young person and their parents or carers, as soon as possible, and:

- include activities that promote physical health and social inclusion, especially education, but also employment, volunteering and other occupations such as leisure activities
- provide support to help the child or young person and their parents or carers realise the plan
- give an up-to-date written copy of the care plan to the young person and their parents or carers if the young person agrees to this; give a copy of the care plan to the parents or carers of younger children; agree a suitable time to review it
- send a copy to the primary healthcare professional who made the referral⁵.

1.3.7 Support children and young people to develop strategies, including risk- and self-management plans, to promote and maintain independence and self-efficacy, wherever possible. Incorporate these strategies into the care plan⁵.

1.3.8 If the child or young person is at risk of crisis, develop a crisis plan with the parents or carers of younger children, or jointly with the young person and their parents or carers, and with their care coordinator. The plan should be respected and implemented, incorporated into the care plan and include:

- possible early warning signs of a crisis and coping strategies
- support available to help prevent hospitalisation
- where the child or young person would like to be admitted in the event of hospitalisation
- definitions of the roles of primary and secondary care professionals and the degree to which parents or carers are involved

- information about 24-hour access to services
- the names of key clinical contacts⁵.

1.3.9 For children and young people with first episode psychosis who are unable to attend mainstream school or college, facilitate alternative educational input in line with their capacity to engage with educational activity and according to their individual needs, with an ultimate goal of returning to mainstream education, training or employment.

1.3.10 If the child or young person and/or their parent or carer is unhappy about the assessment, diagnosis or care plan, give them time to discuss this and offer them the opportunity for a second opinion⁵.

Treatment options for first episode psychosis

1.3.11 For children and young people with first episode psychosis offer:

- oral antipsychotic medication⁷ (see recommendations 1.3.14–1.3.25) in conjunction with
- psychological interventions (family intervention with individual CBT, delivered as set out in recommendations 1.3.26–1.3.32).

1.3.12 If the child or young person and their parents or carers wish to try psychological interventions (family intervention with individual CBT) alone without antipsychotic medication, advise that psychological interventions are more effective when delivered in conjunction with antipsychotic medication. If the child or young person and their parents or carers still wish to try psychological interventions alone, then offer family intervention with individual CBT. Agree a time limit (1 month or less) for reviewing treatment options, including introducing antipsychotic medication. Continue to monitor symptoms, level of distress, impairment and level of functioning, including educational engagement and achievement, regularly.

1.3.13 If the child or young person shows symptoms and behaviour sufficient for a diagnosis of an affective psychosis or disorder, including bipolar disorder and unipolar

⁷ At the time of publication (January 2013), most antipsychotic medication did not have a UK marketing authorisation specifically for children and young people. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. Informed consent should be obtained and documented. See the General Medical Council's [Good practice in prescribing medicines – guidance for doctors](#) for further information.

psychotic depression, follow the recommendations in [Bipolar disorder](#) (NICE clinical guideline 38) or [Depression in children and young people](#) (NICE clinical guideline 28).

Choice of antipsychotic medication

1.3.14 The choice of antipsychotic medication⁷ should be made by the parents or carers of younger children, or jointly with the young person and their parents or carers, and healthcare professionals. Provide age-appropriate information and discuss the likely benefits and possible side effects of each drug including:

- metabolic (including weight gain and diabetes)
- extrapyramidal (including akathisia, dyskinesia and dystonia)
- cardiovascular (including prolonging the QT interval)
- hormonal (including increasing plasma prolactin)
- other (including unpleasant subjective experiences).

How to use oral antipsychotic medication,

1.3.15 Before starting antipsychotic medication⁷, undertake and record the following baseline investigations⁸:

- weight and height (both plotted on a growth chart)
- waist and hip circumference
- pulse and blood pressure
- fasting blood glucose, glycosylated haemoglobin (HbA_{1c}), blood lipid profile and prolactin levels
- assessment of any movement disorders
- assessment of nutritional status, diet and level of physical activity.

1.3.16 Before starting antipsychotic medication, offer the child or young person an electrocardiogram (ECG) if:

- specified in the SPC for adults and/or children

⁸ See [Supplementary information](#) for a table of baseline investigations and monitoring for children and young people who are prescribed antipsychotic medication (read in conjunction with the BNF, BNFC and SPC).

- a physical examination has identified specific cardiovascular risk (such as diagnosis of high blood pressure)
- there is a personal history of cardiovascular disease
- there is a family history of cardiovascular disease such as premature sudden cardiac death or prolonged QT interval, or
- the child or young person is being admitted as an inpatient⁶.

1.3.17 Treatment with antipsychotic medication⁷ should be considered an explicit individual therapeutic trial. Include the following:

- From a discussion with the child or young person and their parent or carer, record the side effects the child or young person is most and least willing to tolerate.
- Record the indications and expected benefits and risks of oral antipsychotic medication, and the expected time for a change in symptoms and appearance of side effects.
- At the start of treatment give a dose below the lower end of the licensed range for adults if the drug is not licensed for children and young people and at the lower end of the licensed range if the drug is licensed for children and young people; slowly titrate upwards within the dose range given in the British national formulary (BNF), the British national formulary for children (BNFC) or the SPC.
- Justify and record reasons for dosages above the range given in the BNF, BNFC or SPC.
- Record the rationale for continuing, changing or stopping medication, and the effects of such changes.
- Carry out a trial of the medication at optimum dosage for 4–6 weeks⁶.

1.3.18 Monitor and record the following regularly and systematically throughout treatment, but especially during titration⁸:

- efficacy, including changes in symptoms and behaviour
- side effects of treatment, taking into account overlap between certain side effects and clinical features of schizophrenia (for example, the overlap between akathisia and agitation or anxiety)

- the emergence of movement disorders
- weight, weekly for the first 6 weeks, then at 12 weeks and then every 6 months (plotted on a growth chart)
- height every 6 months (plotted on a growth chart)
- waist and hip circumference every 6 months (plotted on a percentile chart)
- pulse and blood pressure (plotted on a percentile chart) at 12 weeks and then every 6 months
- fasting blood glucose, HbA_{1c}, blood lipid and prolactin levels at 12 weeks and then every 6 months
- adherence
- physical health.

The secondary care team should maintain responsibility for monitoring physical health and the effects of antipsychotic medication in children and young people for at least the first 12 months or until their condition has stabilised. Thereafter, the responsibility for this monitoring may be transferred to primary care under shared care arrangements.

- 1.3.19 Discuss any non-prescribed therapies that children or young people, or their parents or carers, wish to use (including complementary therapies) with them. Discuss the safety and efficacy of the therapies, and possible interference with the therapeutic effects of prescribed medication and psychological interventions⁶.
- 1.3.20 Discuss the use of alcohol, tobacco, prescription and non-prescription medication and illicit drugs with the child or young person, and their parents or carers where this has been agreed. Discuss their possible interference with the therapeutic effects of prescribed medication and psychological interventions and the potential of illicit drugs to exacerbate psychotic symptoms⁶.
- 1.3.21 'As required' (p.r.n.) prescriptions of antipsychotic medication should be made as described in recommendation 1.3.17. Review clinical indications, frequency of administration, therapeutic benefits and side effects at least weekly. Check whether 'p.r.n.' prescriptions have led to a dosage above the maximum specified in the BNF, BNFC or SPC⁶.

- 1.3.22 Do not use a loading dose of antipsychotic medication (often referred to as 'rapid neuroleptisation')⁶.
- 1.3.23 Do not initiate regular combined antipsychotic medication, except for short periods (for example, when changing medication)⁶.
- 1.3.24 If prescribing chlorpromazine, warn of its potential to cause skin photosensitivity. Advise using sunscreen if necessary⁶.
- 1.3.25 Review antipsychotic medication annually, including observed benefits and any side effects.

How to deliver psychological interventions

- 1.3.26 When delivering psychological interventions for children and young people with psychosis or schizophrenia, take into account their developmental level, emotional maturity and cognitive capacity, including any learning disabilities, sight or hearing problems or delays in language development.
- 1.3.27 Family intervention should:
- include the child or young person with psychosis or schizophrenia if practical
 - be carried out for between 3 months and 1 year
 - include at least 10 planned sessions
 - take account of the whole family's preference for either single-family intervention or multi-family group intervention
 - take account of the relationship between the parent or carer and the child or young person with psychosis or schizophrenia
 - have a specific supportive, educational or treatment function and include negotiated problem solving or crisis management work⁶.

1.3.28 CBT should be delivered on a one-to-one basis over at least 16 planned sessions (although longer may be needed) and:

- follow a treatment manual⁹ so that:
 - children and young people can establish links between their thoughts, feelings or actions and their current or past symptoms, and/or functioning
 - the re-evaluation of the child or young person's perceptions, beliefs or reasoning relates to the target symptoms
- also include at least one of the following components:
 - normalising, leading to understanding and acceptability of their experience
 - children and young people monitoring their own thoughts, feelings or behaviours with respect to their symptoms or recurrence of symptoms
 - promoting alternative ways of coping with the target symptom
 - reducing distress
 - improving functioning⁶.

Monitoring and reviewing psychological interventions

1.3.29 When providing psychological interventions, routinely and systematically monitor a range of outcomes across relevant areas, including the child or young person's satisfaction and, if appropriate, parents' or carers' satisfaction⁶.

1.3.30 Healthcare teams working with children and young people with psychosis or schizophrenia should identify a lead healthcare professional within the team whose responsibility is to monitor and review:

- access to and engagement with psychological interventions
- decisions to offer psychological interventions and equality of access across different ethnic groups⁶.

⁹ Treatment manuals that have evidence for their efficacy from clinical trials are preferred. If developed for adults, the approach should be adapted to suit the age and developmental level of the child or young person.

Competencies for delivering psychological interventions

- 1.3.31 Healthcare professionals delivering psychological interventions should:
- have an appropriate level of competence in delivering the intervention to children and young people with psychosis or schizophrenia
 - be regularly supervised during psychological therapy by a competent therapist and supervisor⁶.
- 1.3.32 Trusts should provide access to training that equips healthcare professionals with the competencies required to deliver the psychological interventions for children and young people recommended in this guideline⁶.

1.4 *Subsequent acute episodes of psychosis or schizophrenia*

- 1.4.1 For children and young people with an acute exacerbation or recurrence of psychosis or schizophrenia offer:
- oral antipsychotic medication⁷ in conjunction with
 - psychological interventions (family intervention with individual CBT).

Pharmacological interventions

- 1.4.2 For children or young people with an acute exacerbation or recurrence of psychosis or schizophrenia, offer oral antipsychotic medication or review existing medication⁷. The choice of drug should be influenced by the same criteria recommended for starting treatment (see recommendations 1.3.14–1.3.25). Take into account the clinical response to and side effects associated with current and previous medication, and monitor as described in recommendation 1.3.18⁶.
- 1.4.3 Aripiprazole is recommended as an option for the treatment of schizophrenia in people aged 15 to 17 years who are intolerant of risperidone, or for whom risperidone is contraindicated, or whose schizophrenia has not been adequately controlled with risperidone. [This recommendation is from [Aripiprazole for the treatment of schizophrenia in people aged 15 to 17 years](#) (NICE technology appraisal guidance 213).]

Psychological and psychosocial interventions

- 1.4.4 Offer family intervention (delivered as set out in recommendation 1.3.27) to all families of children and young people with psychosis or schizophrenia, particularly for preventing and reducing relapse. This can be started either during the acute phase or later, including in inpatient settings⁶.
- 1.4.5 Offer CBT (delivered as set out in recommendation 1.3.28) to all children and young people with psychosis or schizophrenia, particularly for symptom reduction. This can be started either during the acute phase or later, including in inpatient settings⁶.
- 1.4.6 Consider arts therapies (for example, dance movement, drama, music or art therapy) for all children and young people with psychosis or schizophrenia, particularly for the alleviation of negative symptoms. This can be started either during the acute phase or later, including in inpatient settings⁶.
- 1.4.7 If arts therapies are considered, they should be provided by Health Professions Council (HPC) registered arts therapists, with experience of working with children and young people with psychosis or schizophrenia. The intervention should be provided in groups unless difficulties with acceptability and access and engagement indicate otherwise. Arts therapies should combine psychotherapeutic techniques with activity aimed at promoting creative expression, which is often unstructured and led by the child or young person. Aims of arts therapies should include:
- enabling children and young people with psychosis or schizophrenia to experience themselves differently and to develop new ways of relating to others
 - helping children and young people to express themselves and to organise their experience into a satisfying aesthetic form
 - helping children and young people to accept and understand feelings that may have emerged during the creative process (including, in some cases, how they came to have these feelings) at a pace suited to them⁶.
- 1.4.8 Do not routinely offer counselling and supportive psychotherapy (as specific interventions) to children and young people with psychosis or schizophrenia. However, take the child or young person's and their parents' or carers' preferences

into account, especially if other more efficacious psychological interventions, such as CBT, family intervention and arts therapies, are not available locally⁶.

- 1.4.9 Do not offer adherence therapy (as a specific intervention) to children and young people with psychosis or schizophrenia⁶.
- 1.4.10 Do not routinely offer social skills training (as a specific intervention) to children and young people with psychosis or schizophrenia⁶.
- 1.4.11 When psychological interventions, including arts therapies, are started in the acute phase (including in inpatient settings), the full course should be continued after discharge without unnecessary interruption⁶.

1.5 Referral in crisis and challenging behaviour

- 1.5.1 When a child or young person is referred in crisis they should be seen by specialist mental health secondary care services within 4 hours of referral⁵.
- 1.5.2 To avoid admission, aim to:
- explore with the child or young person and their parents or carers what support systems they have, including other family members and friends
 - support a child or young person in crisis and their parents or carers in their home environment
 - make early plans to help the child or young person maintain their day-to-day activities, including education, work, voluntary work, and other occupations and leisure activities, wherever possible⁵.
- 1.5.3 At the end of a crisis assessment, ensure that the decision to start home treatment depends not on the diagnosis, but on:
- the level of distress
 - the severity of the problems
 - the vulnerability of the child or young person and issues of safety and support at home
 - the child or young person's cooperation with treatment⁵.

-
- 1.5.4 Consider the support and care needs of parents or carers of children or young people in crisis. Where needs are identified, ensure they are met when it is safe and practicable to do so⁵.
- 1.5.5 Follow the recommendations in [Self-harm](#) (NICE clinical guideline 16) when managing acts of self-harm in children and young people with psychosis or schizophrenia who are 8 years or over⁶.

Hospital care

- 1.5.6 If a child or young person needs hospital care, this should be in a setting appropriate to their age and developmental level.
- 1.5.7 Before referral for hospital care, think about the impact on the child or young person and their parents, carers and other family members, especially when the inpatient unit is a long way from where they live. Consider alternative care within the community wherever possible. If hospital admission is unavoidable, provide support for parents or carers when the child or young person is admitted.
- 1.5.8 Give verbal and written information to children and young people with psychosis or schizophrenia admitted to hospital, and their parents or carers, about:
- the hospital and the ward in which the child or young person will stay
 - treatments, activities and services available
 - expected contact from health and social care professionals
 - rules of the ward (including substance misuse policy)
 - their rights, responsibilities and freedom to move around the ward and outside
 - meal times
 - visiting arrangements.

Make sure there is enough time for the child or young person and their parents or carers to ask questions⁵.

- 1.5.9 Undertake shared decision-making routinely with children or young people in hospital who are of an appropriate developmental level, emotional maturity and cognitive

capacity, including, whenever possible, those who are subject to the Mental Health Act (1983; amended 1995 and 2007). Include their parents or carers if appropriate⁵.

- 1.5.10 Ensure that children and young people of compulsory school age have access to a full educational programme while in hospital. The programme should meet the National Curriculum, be matched to the child or young person's developmental level and educational attainment, and should take account of their illness and degree of impairment.
- 1.5.11 Ensure that children and young people in hospital continue to have access to a wide range of meaningful and culturally appropriate occupations and activities 7 days per week, and not restricted to 9am to 5pm. These should include creative and leisure activities, exercise, self-care and community access activities (where appropriate). Activities should be facilitated by appropriately trained educational, health or social care professionals⁵.
- 1.5.12 Children and young people receiving community care before hospital admission should be routinely visited while in hospital by the health and social care professionals responsible for their community care⁵.
- 1.5.13 Promote good physical health, including healthy eating, exercise and smoking cessation.

Rapid tranquillisation and restraint

- 1.5.14 Healthcare professionals undertaking rapid tranquillisation and/or restraint in children and young people with psychosis or schizophrenia should be trained and competent in undertaking these procedures in children and young people.
- 1.5.15 Occasionally children and young people with psychosis or schizophrenia pose an immediate risk to themselves or others during an acute episode and may need rapid tranquillisation. Be particularly cautious when considering high-potency antipsychotic medication (such as haloperidol) in children and young people, especially those who have not taken antipsychotic medication before, because of the increased risk of acute dystonic reactions in that age group⁶.

- 1.5.16 After rapid tranquillisation, offer the child or young person the opportunity to discuss their experiences. Provide them with a clear explanation of the decision to use urgent sedation. Record this in their notes⁶.

1.6 *Early post-acute period*

- 1.6.1 In the early period of recovery following an acute episode, reflect upon the episode and its impact with the child or young person and their parents or carers, and make plans for recovery and possible future care.
- 1.6.2 Inform the child or young person and their parents or carers that there is a high risk of relapse if medication is stopped in the 1–2 years following an acute episode⁶.
- 1.6.3 If withdrawing antipsychotic medication, undertake gradually and monitor regularly for signs and symptoms of relapse⁶.
- 1.6.4 After withdrawal from antipsychotic medication, continue monitoring for signs and symptoms of relapse for at least 2 years⁶.

1.7 *Promoting recovery and providing possible future care in primary care*

- 1.7.1 Develop and use practice case registers to monitor the physical and mental health of children and young people with psychosis or schizophrenia in primary care⁶.
- 1.7.2 GPs and other primary healthcare professionals should monitor the physical health of children and young people with psychosis or schizophrenia at least once a year. They should bear in mind that people with schizophrenia are at higher risk of cardiovascular disease than the general population.
- 1.7.3 Identify children and young people with psychosis or schizophrenia who smoke or who have high blood pressure, raised lipid levels or increased waist measurement at the earliest opportunity and monitor for the emergence of cardiovascular disease and diabetes.

-
- 1.7.4 Treat children and young people with psychosis or schizophrenia who have diabetes and/or cardiovascular disease in primary care. Use the appropriate NICE guidance for children and young people where available^{6,10}.
- 1.7.5 Healthcare professionals in secondary care should ensure, as part of the care programme approach (CPA) in England and care and treatment plans in Wales, that children and young people with psychosis or schizophrenia receive physical healthcare from primary care as described in recommendations 1.7.2–1.7.4. Healthcare professionals in secondary care should continue to maintain responsibility for monitoring and managing any side effects of antipsychotic medication⁶.
- 1.7.6 When a child or young person with a diagnosis of psychosis or schizophrenia presents with a suspected relapse (for example, with increased psychotic symptoms or a significant increase in the use of alcohol or other substances) and is still receiving treatment, primary healthcare professionals should refer to the crisis section of the care plan. Consider referral to the key clinician or care coordinator identified in the crisis plan⁶.
- 1.7.7 For a child or young person with psychosis or schizophrenia being cared for in primary care, consider referral to secondary care again if there is:
- poor response to treatment
 - non-adherence to medication
 - intolerable side effects from medication or the child or young person or their parents or carers request a review of side effects
 - the child or young person or their parents or carers request psychological interventions not available in primary care
 - comorbid substance misuse
 - risk to self or others⁶.

¹⁰ See [Type 1 diabetes](#) (NICE clinical guideline 15).

1.8 Promoting recovery and providing possible future care in secondary care

1.8.1 Children and young people with psychosis or schizophrenia who are being treated in an early intervention in psychosis service should have access to that service for up to 3 years (or until their 18th birthday, whichever is longer) whatever the age of onset of psychosis or schizophrenia.

Psychological interventions

1.8.2 Offer family intervention to families of children and young people with psychosis or schizophrenia to promote recovery. Deliver family intervention as described in recommendation 1.3.27⁶.

1.8.3 Consider family intervention particularly for families of children and young people with psychosis or schizophrenia who have:

- recently relapsed or are at risk of relapse
- persisting symptoms⁶.

1.8.4 Offer CBT to assist in promoting recovery in children and young people with persisting positive and negative symptoms and for those in remission. Deliver CBT as described in recommendation 1.3.28⁶.

1.8.5 Consider arts therapies (see recommendation 1.4.7) to assist in promoting recovery, particularly in children and young people with negative symptoms⁶.

Pharmacological interventions

1.8.6 The choice of drug⁷ should be influenced by the same criteria recommended for starting treatment (see recommendations 1.3.14–1.3.25)⁶.

1.8.7 Do not use targeted, intermittent dosage maintenance strategies¹¹ routinely. However, consider them for children and young people with psychosis or schizophrenia who are unwilling to accept a continuous maintenance regimen or if

¹¹ Defined as the use of antipsychotic medication only during periods of incipient relapse or symptom exacerbation rather than continuously.

there is another contraindication to maintenance therapy, such as side-effect sensitivity⁶.

Interventions for children and young people whose illness has not responded adequately to treatment

1.8.8 For children and young people with psychosis or schizophrenia whose illness has not responded adequately to pharmacological or psychological interventions:

- review the diagnosis
- establish that there has been adherence to antipsychotic medication⁷, prescribed at an adequate dose and for the correct duration
- review engagement with and use of psychological interventions and ensure that these have been offered according to this guideline; if family intervention has been undertaken suggest CBT; if CBT has been undertaken suggest family intervention for children and young people in close contact with their families
- consider other causes of non-response, such as comorbid substance misuse (including alcohol), the concurrent use of other prescribed medication or physical illness⁶.

1.8.9 Offer clozapine¹² to children and young people with schizophrenia whose illness has not responded adequately to pharmacological treatment despite the sequential use of adequate doses of at least two different antipsychotic drugs each used for 6–8 weeks⁶.

1.8.10 For children and young people whose illness has not responded adequately to clozapine¹² at an optimised dose, consider a multidisciplinary review, and recommendation 1.8.8 (including measuring therapeutic drug levels) before adding a second antipsychotic to augment treatment with clozapine. An adequate trial of such an augmentation may need to be up to 8–10 weeks. Choose a drug that does not compound the common side effects of clozapine⁶.

¹² At the time of publication (January 2013), clozapine did not have a UK marketing authorisation for this indication. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. Informed consent should be obtained and documented. See the General Medical Council's [Good practice in prescribing medicines – guidance for doctors](#) for further information.

Education, employment and occupational activities for children and young people with psychosis and schizophrenia

- 1.8.11 For children and young people of compulsory school age, liaise with the child or young person's school and educational authority, subject to consent, to ensure that ongoing education is provided.
- 1.8.12 Liaise with the child or young person's school and with their parents or carers, subject to consent, to determine whether a special educational needs assessment is necessary. If it is agreed that this is needed, explain to parents or carers how to apply for an assessment and offer support throughout the process.
- 1.8.13 Provide supported employment programmes for those young people with psychosis or schizophrenia above compulsory school age who wish to return to work or find employment. Consider other work-related activities and programmes when individuals are unable to work or are unsuccessful in their attempts to find employment⁶.
- 1.8.14 Mental health services should work in partnership with local stakeholders, including those representing black and minority ethnic groups, to enable young people with psychosis or schizophrenia to access local employment and educational opportunities. This should be sensitive to the young person's needs and skill level and is likely to involve working with agencies such as Jobcentre Plus, disability employment advisers and non-statutory providers⁶.
- 1.8.15 Routinely record the daytime activities of children and young people with psychosis or schizophrenia in their care plans, including educational and occupational outcomes⁶.

2 Research recommendations

The Guideline Development Group has made the following recommendations for research, based on its review of evidence, to improve NICE guidance and patient care in the future.

2.1 *What are the long-term outcomes, both psychotic and non-psychotic, for children and young people with attenuated or transient psychotic symptoms suggestive of a developing psychosis, and can the criteria for 'at risk states' be refined to better predict those who will and those who will not go on to develop psychosis?*

The suggested programme of research would be in two phases. First, a systematic review and meta-analysis of prospective observational studies/cohorts of children and young people identified at high or ultra-high risk of developing psychosis would be undertaken. The review would identify risk and protective factors most strongly associated with the later development of psychotic and non-psychotic outcomes. Second, the factors identified in the first phase would be used to identify a large cohort of children and young people with these factors and to evaluate the effectiveness of these refined criteria for predicting the later development of psychotic and non-psychotic outcomes.

Why this is important

A major problem with trials of treatments for populations of children and young people deemed to be 'at risk' or 'at ultra-high risk' of developing psychosis is identifying the precise symptoms and/or behaviours or (risk) factors that are most strongly associated with the development of psychosis; and conversely, which (protective) factors are likely to be associated with a lowered risk of later psychosis. At present, identified factors have a low predictive value, with only about 10–20% of children and young people who have been identified as at high risk going on to develop psychosis. If these risk and protective factors could be refined, it would be possible to better target children and young people who are most at risk, and reduce the numbers of those thought to be 'at risk' who do not go on to later develop psychosis.

2.2 *What is the clinical and cost effectiveness of omega-3 fatty acids in the treatment of children and young people considered to be at high risk of developing psychosis?*

The suggested programme of research would need to test out, using an adequately powered, multicentre randomised controlled design, the likely benefits and costs of using omega-3 fatty acids for children and young people at high risk of developing psychosis. The outcomes considered should include transition to psychosis, quality of life, symptomatic and functional improvements, treatment acceptability, side effects and self-harm. There should be follow-up at 3 years. The trial should also estimate the cost effectiveness of intervening.

Why this is important

A number of interventions have been trialled in an attempt to avert the development of psychosis, including drugs, psychological interventions and other interventions. A relatively recent, moderate-sized randomised controlled trial of omega-3 fatty acids has shown the best evidence of any intervention, to date, reducing the rates of transition from 'high risk' states to a sustained psychosis. However, this is a single trial, which is underpowered, undertaken in one centre and lacks any health economic analysis.

2.3 *What is the clinical and cost effectiveness for family intervention combined with individual CBT in the treatment of children and young people considered to be at high risk of developing psychosis and their parents or carers?*

The suggested programme of research would need to test out, using an adequately powered, multicentre, randomised controlled design, the likely benefits and costs of providing family intervention, combined with individual CBT, for children and young people at high risk of developing psychosis and their parents or carers. The outcomes considered should include transition to psychosis, quality of life, symptomatic and functional improvements, treatment acceptability and self-harm. There should be follow-up at 3 years. The trial should also estimate the cost effectiveness of intervening.

Why this is important

A number of interventions have been trialled in an attempt to avert the development of psychosis, including drugs, psychological interventions and other interventions. After the first episode of psychosis, family intervention as an adjunct to antipsychotic medication substantially and significantly reduces relapse rates. A single small trial combining CBT family treatment with individual CBT without antipsychotic treatment suggested an important reduction in transition rates to the first psychosis.

2.4 *What is the clinical and cost effectiveness of psychological intervention alone, compared with antipsychotic medication and compared with psychological intervention and antipsychotic medication combined, in young people with first episode psychosis?*

The programme of research would compare the clinical and cost effectiveness of psychological intervention alone, compared with antipsychotic medication, and compared with psychological intervention and antipsychotic medication combined, for young people in the early stages of psychosis using an adequately powered study with a randomised controlled design. The combination of psychological interventions most likely to have an impact is family intervention and individual CBT. The key outcomes should include symptoms, relapse rates, quality of life, treatment acceptability, experience of care, level of psychosocial functioning and the cost effectiveness of the interventions.

Why this is important

The personal and financial cost of psychosis and schizophrenia to the person, their family and friends, and to society is considerable. The personal cost is reflected in a suicide rate of nearly 15% among people with schizophrenia, a lifelong unemployment rate that varies between 50 and 75%, depending on geographical location, and reduced life expectancy. The additional cost to the healthcare system for one person with schizophrenia is estimated to reach over £50,000 per year, on average, throughout their life.

Currently, the mainstay of treatment is antipsychotic medication, but the potential adverse effects are such that there is considerable impetus to develop alternative treatment strategies to

allow either lower doses or to remove the need for medication entirely. It has been recognised that psychological interventions as an adjunct to antipsychotic medication have an important part to play in the treatment of schizophrenia. NICE clinical guideline 82 identified family intervention and CBT as adjunct treatments and current evidence suggests that these interventions are cost saving. However, evidence for adjunctive family intervention and CBT is lacking in children and young people with psychosis. Furthermore, there has been one recent positive trial of CBT as a first-line treatment, without antipsychotics, for young people in the early stages of psychosis.

2.5 *What is the clinical effectiveness of clozapine for children and young people with schizophrenia with symptoms unresponsive to antipsychotic medication and psychological treatment combined?*

The suggested programme of research would need to test out, using an adequately powered, randomised controlled design, the likely benefits of using clozapine, compared with another antipsychotic, for children and young people with symptoms of schizophrenia unresponsive to antipsychotic medication and psychological treatment combined. The outcomes considered should include quality of life, symptomatic and functional improvements, treatment acceptability, side effects and length of hospitalisation.

Why this is important

Currently, about 30% of people with schizophrenia have symptoms that do not respond adequately to treatment with an antipsychotic. Although precise figures are unavailable, especially for children and young people, smaller percentages of people do not respond when a second, alternative, antipsychotic and an adequate course of psychological treatment have been tried. For these people, clozapine, which has a different dopamine receptor subtype blocking profile from other antipsychotics, has become an important treatment option in adults. However, evidence is lacking (only one study) about the effectiveness of clozapine for 'treatment-resistant schizophrenia' in children and young people.

2.6 *What is the most effective management strategy for preventing the development of excessive weight gain and metabolic syndrome*

associated with the use of antipsychotic medication in children and young people?

The suggested programme of research would be in two parts: (1) a longitudinal cohort study (a national observational database of at least 12 months' duration) to determine the incidence and predictors of adverse physical effects of antipsychotic medication; (2) a randomised controlled trial of behavioural and/or medical approaches to reduce weight gain and the risk of metabolic syndrome associated with antipsychotic medication.

Why this is important

Rapid weight gain associated with antipsychotic medication and poor physical health (smoking, lack of exercise) leading to type 2 diabetes and metabolic syndrome are major sources of morbidity and premature mortality in young people with psychosis and schizophrenia. Most evidence of adverse effects comes from short-term studies of antipsychotics (maximum 8–12 weeks). In contrast, very little is known about the longer term adverse effects of these drugs. Evidence is needed both on longer term adverse effects as well as on effective early intervention strategies that reduce these risk factors and improve physical health outcomes.

3 Other information

3.1 *Scope and how this guideline was developed*

NICE guidelines are developed in accordance with a [scope](#) that defines what the guideline will and will not cover.

How this guideline was developed

NICE commissioned the National Collaborating Centre for Mental Health to develop this guideline. The Centre established a Guideline Development Group (see [section 4](#)), which reviewed the evidence and developed the recommendations.

The methods and processes for developing NICE clinical guidelines are described in [The guidelines manual](#).

3.2 *Related NICE guidance*

Details are correct at the time of publication (January 2013). Further information is available on the [NICE website](#).

Published

General

- [Service user experience in adult mental health](#). NICE clinical guidance 136 (2011).
- [Medicines adherence](#). NICE clinical guideline 76 (2009).

Condition-specific

- [Aripiprazole for the treatment of schizophrenia in people aged 15 to 17 years](#). NICE technology appraisal 213 (2011).
- [Alcohol-use disorders](#). NICE clinical guideline 115 (2011).
- [Schizophrenia](#). NICE clinical guideline 82 (2009).
- [Borderline personality disorder](#). NICE clinical guideline 78 (2009).
- [Antisocial personality disorder](#). NICE clinical guideline 77 (2009).
- [Drug misuse: opioid detoxification](#). NICE clinical guideline 52 (2007).
- [Drug misuse: psychosocial interventions](#). NICE clinical guideline 51 (2007).
- [Obesity](#). NICE clinical guideline 43 (2006).
- [Bipolar disorder](#). NICE clinical guideline 38 (2006).
- [Obsessive-compulsive disorder](#). NICE clinical guideline 31 (2005).
- [Depression in children and young people](#). NICE clinical guideline 28 (2005).
- [Post-traumatic stress disorder](#). NICE clinical guideline 26 (2005).
- [Violence](#). NICE clinical guideline 25 (2005).
- [Self-harm](#). NICE clinical guideline 16 (2004).
- [Type 1 diabetes](#). NICE clinical guideline 15 (2004).

Under development

NICE is developing the following guidance (details available from the [NICE website](#)):

- Bipolar disorder (update). NICE clinical guideline. Publication date to be confirmed.

-
- Diabetes in children (update). NICE clinical guideline. Publication date to be confirmed.
 - Social anxiety disorder. NICE clinical guideline. Publication date to be confirmed.

4 The Guideline Development Group, National Collaborating Centre and NICE project team

Guideline Development Group

Chris Hollis (Chair, Guideline Development Group)

Professor of Child and Adolescent Psychiatry, University of Nottingham

Tim Kendall (Facilitator, Guideline Development Group)

Director, National Collaborating Centre for Mental Health, Royal College of Psychiatrists; Medical Director/Consultant Adult Psychiatrist, Sheffield Health and Social Care NHS Foundation Trust; Visiting Professor, University College London

Max Birchwood

Professor of Youth Mental Health, University of Birmingham; Clinical Director, YouthSpace Mental Health Programme, Birmingham and Solihull Mental Health NHS Foundation Trust

Rory Byrne

Service User Representative; Researcher, Greater Manchester West Mental Health NHS Foundation Trust

Andrew Clark

Consultant in Adolescent Psychiatry, Greater Manchester West Mental Health NHS Foundation Trust

Jaeta Egoh

Service User Representative

Elena Garralda

Professor and Honorary Consultant in Child and Adolescent Psychiatry, Imperial College London and Central and North West London NHS Foundation Trust

Laura Graham

Carer Representative and Involvement Worker and Young Person's Panel Advisor for Rethink Mental Illness

Anthony James

Consultant Child and Adolescent Psychiatrist and Honorary Senior Lecturer, Oxfordshire and Buckinghamshire Mental Health NHS Foundation Trust

Tim McDougall

Nurse Consultant, Clinical Director (Tier 4 CAMHS) and Lead Nurse (CAMHS), Cheshire and Wirral Partnership NHS Foundation Trust

Anthony Morrison

Professor of Clinical Psychology, Greater Manchester West Mental Health NHS Foundation Trust

Gillian Rose

Consultant Child and Adolescent Psychiatrist, Central and North West London NHS Foundation Trust

David Shiers

GP Advisor to the National Audit of Schizophrenia (The Royal College of Psychiatrists) and Rethink Mental Illness Trustee

Kirsty Smedley

Consultant Clinical Psychologist, The Priory Hospital Cheadle Royal; Honorary Lecturer, Manchester University

Darryl Thompson

Psychosocial Interventions Development Lead, South West Yorkshire Partnership NHS Foundation Trust

David Ward

Consultant Adolescent Psychiatrist, Newcastle CAMHS and Early Intervention in Psychosis Service (2010–2011)

National Collaborating Centre for Mental Health

Henna Bhatti

Research Assistant (2010–2011)

Melissa Chan

Systematic Reviewer (2010–2011)

Nadir Cheema

Health Economist

Marie Halton

Research Assistant (2010–2011)

Hannah Jackson

Research Assistant (2011–2012)

Linnéa Larsson

Project Manager and Research Assistant (2012–2013)

Christina Loucas

Research Assistant (2012–2013)

Kate Satrettin

Project Manager (2010–2012)

Christine Sealey

Associate Director (Operations)

Megan Stafford

Systematic Reviewer (2011–2013)

Sarah Stockton

Senior Information Scientist

Clare Taylor

Senior Editor

NICE project team

Sarah Willett

Associate Director

Caroline Keir

Guideline Commissioning Manager

Margaret Ghلامي

Guideline Coordinator (2012–2013)

Jennifer Heaton

Guideline Coordinator (2010–2012)

Judith Thornton

Technical Lead

Jasdeep Hayre

Health Economist

Anne-Louise Clayton

Editor

Supplementary information on baseline investigations and monitoring

Table 1 Baseline investigations and monitoring for children and young people who are prescribed antipsychotic medication (read in conjunction with the BNF, BNFC and SPC)

	Baseline investigations before starting antipsychotic medication	Monitor weekly for the first 6 weeks	Monitor at 12 weeks	Monitor every 6 months thereafter	Monitor regularly throughout treatment, and especially during titration
Weight ¹ (plotted on a growth chart)	Yes	Yes	Yes	Yes	
Height ¹ (plotted on a growth chart)	Yes			Yes	
Waist and hip circumference (plotted on a percentile chart)	Yes			Yes	
Pulse	Yes		Yes	Yes	
Blood pressure (plotted on a percentile chart)	Yes		Yes	Yes	
Fasting blood glucose	Yes		Yes	Yes	
HbA _{1c} (glycosylated haemoglobin)	Yes		Yes	Yes	
Blood lipid profile	Yes		Yes	Yes	
Prolactin level	Yes		Yes	Yes	
Movement disorders (extrapyramidal symptoms, akathisia, dystonia and tardive dyskinesia)	Yes				Yes ²

	Baseline investigations before starting antipsychotic medication	Monitor weekly for the first 6 weeks	Monitor at 12 weeks	Monitor every 6 months thereafter	Monitor regularly throughout treatment, and especially during titration
Nutritional status, diet and level of physical activity	Yes				Yes
The side effects the child or young person is most or least willing to tolerate	Yes				
ECG	Yes ³				
Efficacy					Yes
Side effects					Yes
Adherence					Yes
<p>¹ Calculate and document BMI (percentile).</p> <p>² Even if no baseline assessment (and at each clinic visit if more frequent).</p> <p>³ If specified in the SPC for adults and/or children; a physical examination has identified specific cardiovascular risk (such as diagnosis of high blood pressure); there is personal history of cardiovascular disease; there is a family history of cardiovascular disease such as sudden cardiac death or prolonged QT interval; or the child or young person is being admitted as an inpatient.</p>					

About this guideline

NICE clinical guidelines are recommendations about the treatment and care of people with specific diseases and conditions in the NHS in England and Wales.

NICE guidelines are developed in accordance with a scope that defines what the guideline will and will not cover.

This guideline was developed by the National Collaborating Centre for Mental Health, which is based at the Royal College of Psychiatrists. The Collaborating Centre worked with a Guideline Development Group, comprising healthcare professionals (including consultants, GPs and

nurses), patients and carers, and technical staff, which reviewed the evidence and drafted the recommendations. The recommendations were finalised after public consultation.

The methods and processes for developing NICE clinical guidelines are described in [The guidelines manual](#).

Strength of recommendations

Some recommendations can be made with more certainty than others. The Guideline Development Group makes a recommendation based on the trade-off between the benefits and harms of an intervention, taking into account the quality of the underpinning evidence. For some interventions, the Guideline Development Group is confident that, given the information it has looked at, most patients would choose the intervention. The wording used in the recommendations in this guideline denotes the certainty with which the recommendation is made (the strength of the recommendation).

For all recommendations, NICE expects that there is discussion with the patient about the risks and benefits of the interventions, and their values and preferences. This discussion aims to help them to reach a fully informed decision (see also [Person-centred care](#)).

Interventions that must (or must not) be used

We usually use 'must' or 'must not' only if there is a legal duty to apply the recommendation. Occasionally we use 'must' (or 'must not') if the consequences of not following the recommendation could be extremely serious or potentially life threatening.

Interventions that should (or should not) be used – a 'strong' recommendation

We use 'offer' (and similar words such as 'refer' or 'advise') when we are confident that, for the vast majority of patients, an intervention will do more good than harm, and be cost effective. We use similar forms of words (for example, 'Do not offer...') when we are confident that an intervention will not be of benefit for most patients.

Interventions that could be used

We use 'consider' when we are confident that an intervention will do more good than harm for most patients, and be cost effective, but other options may be similarly cost effective. The choice of intervention, and whether or not to have the intervention at all, is more likely to depend

on the patient's values and preferences than for a strong recommendation, and so the healthcare professional should spend more time considering and discussing the options with the patient.

Recommendation wording in adapted recommendations

NICE began using this approach to denote the strength of recommendations in guidelines that started development after publication of the 2009 version of [The guidelines manual](#) (January 2009). This does not apply to any recommendations that have been adapted from guidelines that started development before this. In particular, in adapted recommendations the word 'consider' may not necessarily be used to denote the strength of the recommendation.

Other versions of this guideline

The full guideline, [Psychosis and schizophrenia in children and young people: recognition and management](#) contains details of the methods and evidence used to develop the guideline. It is published by the National Collaborating Centre for Mental Health.

The recommendations from this guideline have been incorporated into a [NICE Pathway](#).

We have produced [information for the public](#) about this guideline.

Implementation

[Implementation tools and resources](#) to help you put the guideline into practice are also available.

Your responsibility

This guidance represents the view of NICE, which was arrived at after careful consideration of the evidence available. Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. However, the guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer, and informed by the summaries of product characteristics of any drugs.

Implementation of this guidance is the responsibility of local commissioners and/or providers. Commissioners and providers are reminded that it is their responsibility to implement the

guidance, in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity and foster good relations. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with those duties.

Copyright

© National Institute for Health and Clinical Excellence 2013. All rights reserved. NICE copyright material can be downloaded for private research and study, and may be reproduced for educational and not-for-profit purposes. No reproduction by or for commercial organisations, or for commercial purposes, is allowed without the written permission of NICE.

Contact NICE

National Institute for Health and Clinical Excellence
Level 1A, City Tower, Piccadilly Plaza, Manchester M1 4BT

www.nice.org.uk

nice@nice.org.uk

0845 033 7780